BENIGN PROSTATIC HYPERPLASIA (BPH)

TREATMENT OPTIONS



When the symptoms of BPH cannot be controlled through conventional treatment, several interventions now exist. To help you make an informed decision, below is a list of these options, as well as the pros and cons of each approach.

Procedure	Treatment Description	Requires hospitalization and recovery?	Requires general anesthesia?	Requires permanent implants?	Preserves sexual function?	Suitable for larger prostate?	Incidence of side effects	Need for re-treatment	Common side effects, in addition to normal surgical or procedure risk of infection, include:
Prostatectomy	Recommended for those with a very enlarged prostate, this open surgery procedure is used to remove a large amount of prostate tissue or gland itself.	/	/	×	×	/	High	Low	Stress incontinence Urinary tract infection Erectile dysfunction Dry orgasm
Transurethral Resection of the Prostate (TURP)	Common surgery to treat BPH, this approach cuts and removes excess prostate tissue to make gland smaller and reduce symptoms.			X	X		High	Low	Retrograde ejaculation Erectile dysfunction Dry orgasm Loss of bladder control
Transurethral Incision of the Prostate (TUIP)	Surgery involves making small cuts within the prostate gland to reduce pressure on urethra.	/	/	×	X	X	Medium	High	Temporary difficulty urinating Urinary tract infection Need for re-treatment
UroLift [®]	Non-invasive procedure that uses a special device to place permanent implants into body to lift prostate tissue away from urethra.	X	X	/	/	X	Low	High	Painful urination Urinary urgency Light blood in urine
Transurethral Microwave Therapy (TUMT)	Non-invasive procedure that uses a microwave antenna on a tube inserted into bladder. The microwave heat destroys excess prostate tissue.	×	×	×	/	×	Medium	High	New or worsening urinary symptoms Chronic prostate inflammation Temporary painful urination
Rezum Water Vapor Therapy	Device inserted into urethra delivers water vapor to treat excess prostate tissue.	×	×	×	/	×	Low	Low	Temporary painful urination Urinary urgency Light blood in urine Decrease in ejaculatory volume
Prostate Artery Embolization	Non-invasive procedure to partially block blood flow to prostate gland. This causes the gland to shrink and symptoms to alleviate. Also appropriate for those with larger prostates.	×	×	×	/	/	Low	Low	Temporary painful urination Light blood in urine Nausea